

Particle Signatures

Fermilab 2009



The ArgoNeuT LArTPC: a dedicated Experiment for neutrino Cross Section measurement at FNAL

NOW 2010

September 7th, 2010

Ornella Palamara
Laboratori Nazionali del Gran Sasso


✓ In the recent years, due to the increasing interest on LAr-TPC technology in the US, a dedicated experiment (ArgoNeuT, 2007) has been included as a first step in a graded program towards massive LBL neutrino oscillation experiments.

✓ One of the main uncertainties in the next generation long baseline oscillation experiments is given by the neutrino-nucleus interaction cross section in the “few-GeV region”.

The (CC-QE) Cross Section (how well we measured it)

• Charged-Current Quasi-Elastic Scattering

Second generation measurements



- MiniBooNE/SciBooNE in agreement, but tension with higher energy NOMAD results. All three on carbon. This is not understood.
- Single point, first and so far unique investigation with Ar target, in agreement with NOMAD data (same, *high energy* ν beam - WANF)

ArgoNeuT Physics Goals




- Measure charged-current cross-section in the “few GeV” (1-5 GeV) range:
 - CC Quasi-Elastic (QE) channel
 - CC Resonant (RES: $\Delta \rightarrow \pi N$) channelwith unprecedented sensitivity to **products of FSI** (vertex activity characterization)
- e/γ separation study and optimization \Rightarrow superior background rejection
 - Important for ν_e appearance: excellent signal (CC ν_e) efficiency and background (NC π^0) rejection
 - Particle identification from energy deposition (dE/dx) measured along track
- Develop reconstruction techniques useful for all future LArTPCs:
 - full 3D reconstruction of the event topology
 - precise calorimetric reconstruction of deposited energy and Particle Identification



ArgoNeuT



- ✓ **ArgoNeuT** is a 175 liter (active) Liquid Argon Time Projection Chamber (LArTPC)
- ✓ Jointly funded by DOE/NSF
- ✓ Designed and **assembled in 2007-08**, first **commissioned** (on surface) at FNAL in **Summer 2008**
- ✓ Moved underground in the **NuMI beam** at FNAL, in front of **MINOS Near Detector**, **early 2009**
- ✓ **Phase I**: Exposure to $\nu/\bar{\nu}$ beam (*LE beam option*): **June'09 \oplus Sept'09-Feb.'10**
- ✓ **Phase II - (second run)**: in the *BOOSTER low-energy nu-beam (SciBooNE enclosure)* - **2011**




Fermilab, NuMI beam line



MINOS Hall: ArgoNeuT just upstream of the MINOS ND


ArgoNeuT Design



The TPC, about to enter the inner cryostat

2 read-out planes: **Induction and Collection**
each channel: 2048 samples in 400 microseconds

Cryostat Volume	500 Liters
TPC Volume	175 Liters
# Electronic Channels	480
Wire Pitch	4 mm
Electronics Style (Temperature)	JFET (293 K)
Max. Drift Length (Time)	0.5m (330 μ s)
Electric field	500 V/cm




The fully-instrumented detector in the beamline

ArgoNeuT's physics run in the NuMI beam





Schematic of NuMI experiments




NuMI beam Fluxes - Low Energy (LE) mode

$\langle E_\nu \rangle = 3.7 \text{ GeV}$



ArgoNeuT POT delivered and accumulated



Reaction	#events in ($\sim 1.35\text{E}20$ POT)
ν_μ CC	~ 6600
$\bar{\nu}_\mu$ CC	~ 4900
ν_μ CCQE	~ 600
ν_e CC	~ 130

- ✓ Stable, shift-free operation for >5 months!
- ✓ The first 1000s of (anti-)neutrino LArTPC events collected in a low-energy ($\sim 3 \text{ GeV}$) neutrino beam ever!


(Neutrino) Event Display




The detector provides two 2D-views of the event.

The color scale is indicative of the energy deposited along the track.

- m.i.p. yield: ~ 6000 e/mm
- Very fine pixel size (4mmx4mm x 0.3mm)
- Dark “shadow bands” are due to electronics returning to baseline...
- Fourier decomposition (FFT) to remove electronics response (Filtering).




Wire pulse



Neutrino Event – ν_e



El.m. shower (not fully contained) +
short densely ionizing track at the vertex




✓ This (beam-intrinsic) event demonstrates what a signal-like electron-neutrino event looks like to in LArTPC.

✓ Current and future long baseline neutrino oscillation experiments (MINOS, T2K, NoVA, LBNE, ...) search for electron-neutrino appearance in order to measure θ_{13} and δ_{CP} .


ν event Reconstruction

Offline reconstruction procedure:


1. Hit identification
2. Hit reconstruction
3. Cluster/Vertex reconstruction
4. 3D track reconstruction



Hit finding + density-based clustering.



3D reconstruction



Track finding/fitting + vertex/endpoint finding


5. Matching ArgoNeuT tracks with downstream MINOS ND

for escaping muon momentum reconstruction and sign determination


6. Calorimetric reconstruction

7. Particle Identification (dE/dx along the track)

μ from upstream ν beam interaction




3D reconstruction




(many) crossing μ 's superimposed in 3D view


Muon calorimetric reconstruction



Muon angular distribution





μ from upstream $\bar{\nu}$ beam interaction: Matching with MINOS ND (I)




Tracks whose direction extrapolated from ArgoNeuT matches a MINOS track

Difference between horizontal coordinates and vertical coordinates of the “matched tracks”





μ from upstream $\bar{\nu}$ beam interaction: Matching with MINOS ND (II)



MINOS:
reconstruction of the
“matched muon” energy

Muon momentum reconstruction from MINOS ND:
- by curvature in magn. field - 12% resolution for a 10 GeV muon
- by range for stopping muons ~6% resolution)



MINOS:
measurement of the
“matched muon” sign



Particle ID




Penetrating particle
(from upstream interaction)
stopping in LAr volume



Minimum ionizing ptcl:
muon or pion


Track length= 52 cm
Kinetic Energy=160 MeV
(in agreement with expectations
GEANT)

Evolution of the ionization along the track



Muon-Pion separation possible only in some cases

Particle ID



Muon-Pion separation possible only in same cases

ν_μ CC QE event reconstruction



ν Interaction in LAr volume

$\mu+p$ (ν_μ CC QE event)

+



*uncorrelated tracks from
upstream neutrino interaction*

Heavy ionizing ptcl.



m.i.p. particle





Muon






ν interaction in LAr

$\mu + \pi$ event + proton track



2 m.i.p. particles ($\mu + \pi$) at vtx
[faking a QEL signature ($\mu + p$)]


+
proton track (far from vtx.)




PID: e/γ separation study and optimization

- Photon conversion background to ν_e interactions
 - Separation from primary vertex or by double ionization
 - γ -conversion over a minimum ionizing track requires excellent pair resolution

Careful inspection yields a minimum ionizing track with overlapping γ conversion





Energy loss in the first 24mm of track: 250 MeV electrons vs. 250 MeV gammas



dE/dx for electrons and gammas in first 2.4 cm of track

$$\pi^0 \rightarrow \gamma \gamma \rightarrow (e^+e^-) (e^+e^-)$$





For e^+e^- efficiency $> 80\%$,
 e contamination $< 5\%$

Understanding vertex activity

“Final State (re-)Interactions”- *the main source of uncertainty:*



even the “easiest” topology (CC-QE) is not so simple



FSI

- **proton Spallation** (intranuclear interactions with **p** and **n** emission ...but also π^\pm, π^0)
- **Nuclear evaporation** (lower kin.en. **p** and **n**)
- (and/or) **Fission** (nuclear fragments, α 's,...)
- **Nuclear de-excitation** with γ emission

These products are usually neglected because not detectable, unless...
.... a high quality imaging detector is in use !!





A zoomed-in view of a CCQE-like neutrino event with evidence of vertex activity



Understanding vertex activity

“Final State (re)-Interactions”- *the main source of uncertainty:*

even the “easiest” topology (CC-QE) is not so simple



FSI



These products are usually neglected because not detectable, unless...
.... a high quality imaging detector is in use !!

A zoomed-in view of a CCQE-like neutrino event with evidence of vertex activity

ArgoNeuT Collaboration

F. Cavanna
University of L'Aquila

A. Ereditato, S. Haug, B. Rossi, M. Weber
University of Bern

B. Baller, C. James, S. Pordes, G. Rameika, B. Rebel
Fermi National Accelerator Laboratory

M. Antonello, O. Palamara
Gran Sasso National Laboratory

T. Bolton, S. Farooq, G. Horton-Smith, D. McKee
Kansas State University

C. Bromberg, D. Edmunds, P. Laurens, B. Page
Michigan State University

K. Lang, R. Mehdiyev
The University of Texas at Austin

C. Anderson, E. Church, B. Fleming, R. Guenette, S. Linden,
K. Partyka, M. Soderberg*, J. Spitz
Yale University



Conclusions



Next-generation neutrino physics experiments require precision

Particle IDentification and
fine grained 3D imaging on very large scale.

Liquid Argon TPC combines an ideal detection medium with a modern imaging and calorimetric readout technique, scalable to very large volume/mass.

ArgoNeuT is a fully operational LArTPC: during the (first) ν -run, large samples of neutrino/antineutrino events have been collected for the 1st time ever in a low-Energy beam.

The extension to a second run period is being proposed at FNAL

Extensive Real data/experience is invaluable in improving LArTPC technique.
Analysis software is being developed as general purpose tool for future LArTPCs.
Highly sophisticated/detailed MonteCarlo codes are needed,
and are currently under test/optimization